

# Sexual Assault and the Brain: Key Info for Investigators, Advocates and Other Professionals

Jim Hopper, Ph.D., August 2016

## Brain-based Effects: Vulnerability and Needs

- Whether reporting a recent sexual assault or one from long ago, the person is very vulnerable.
- They may be tormented by memories and reminders, emotionally 'shut down' and 'numbed out', or cycling between these extremes. **Be careful not to judge credibility based on emotional state.**
- Many symptoms and problems are **attempts to cope**. These include using substances – which may be attempts to escape terrible memories, anxiety, etc. – and compulsive or risky sexual behaviors, which may be attempts to gain a sense of mastery and control over one's sexual experiences.
- Having to talk about the assault can **feel like having one's 'defenses' battered down**. This can cause difficulties in recalling – even when sincerely trying – parts of the assault experience that are particularly disturbing, or about which one feels ashamed. Or, after disclosing such things, they may feel like they did during the assault: violated, overwhelmed and re-traumatized.
- **Most important needs: safety, control, trust, understanding, and compassion.** Find ways to meet these needs within the boundaries of your role, including allowing them to recount what they remember first as an interrupted narrative, then asking (non-leading) follow-up questions. Even simple options and choices, like whether they want a drink, or when to take breaks, can help a lot – improving cooperation and results.

## Brain-based Effects: Fear Circuitry in Control, Prefrontal Cortex Impaired, Running on Habits and Reflexes

- If someone is being sexually assaulted, as long as the person is conscious, even if intoxicated, at some point **the fear circuitry will detect the attack and immediately dominate brain functioning**.
- Within seconds of the fear circuitry taking control, the **prefrontal cortex is impaired**. This results in...
- **Bottom-up attention**: the fear circuitry, not the prefrontal cortex, controls what gets attention.
- **Impairment of prefrontal cortex capacities** for rational thinking, for planning effective responses, for remembering important information (e.g., there are people nearby who would hear a scream), etc.
- **Reflex responses** that are hard-wired into human brains because we evolved as prey, not just predators. These range from a brief '**freeze response**' when attack is detected (in which movement ceases and the brain assesses the attack and possible escape options), to **extreme survival reflexes** including **dissociation** (awareness is disconnected from emotions and body sensations, and one may go on 'autopilot,' including engaging in sex acts), **tonic immobility** (literally unable to move or speak, and rigid body, which is different from freezing), and **collapsed immobility** (muscles are limp, may be 'sleepy,' 'dizzy,' or even pass out).
- **Habit responses** that are rooted in social conditioning, e.g., **how girls and women are socialized** to respond to males' unwanted sexual advances (in nice, polite, face-saving ways), or **habits of dealing with aggressive and dominant people**. Some habit responses were **learned to cope with childhood abuse**.

## Brain-based Effects: Memories

- **Central details**: What the fear circuitry focused attention on during the assault. These tend to be very well encoded into memory, and are **likely to be accurate, consistent and corroborated** (even by perpetrator). They may (at first) not seem central to the investigation (e.g., detailed description of a table or plant), but may be consistent with states of fear and trauma, evidence of being in the described location, etc.
- **Peripheral details**: Details that *did not get attention*, likely because the fear circuitry didn't see them as relevant to survival. They are usually not encoded into memory, or very poorly encoded, therefore **likely to be recalled poorly and/or inconsistently over time**. They may be a central focus of an investigation (e.g., things perpetrator did), but 'failure' to recall such things *does not indicate lack of credibility*; it just means they weren't encoded in the first place, as expected from brains responding to assault, fear and trauma.
- **Contextual information** (e.g., the layout of a room) and **time-sequence information** (e.g., the order in which sexual acts occurred) are usually poorly encoded. Again, an expected impact of 'the brain on fear.'
- Experiences around the time '**when the fear kicked in**' are usually well encoded. Attention is still required for encoding into memory, but because the hippocampus *temporarily* goes into a **super-encoding mode**, memories of when the fear kicked in may include substantial contextual and time-sequence information.